

LATITUDINAL GRADIENT FACILITATES ADAPTIVE POPULATION DIFFERENTIATION
OF *CAMPANULASTRUM AMERICANUM* (AMERICAN BELLFLOWER)

Marisa L. Naciuk

Columbus State University

The College of Letters and Sciences

The Graduate Program in Natural Sciences (Biology Track)

Latitudinal Gradient Facilitates Adaptive Population Differentiation of

Campanulastrum americanum (American Bellflower)

A Thesis in

Natural Science

By

Marisa L. Naciuk

Submitted in Partial Fulfillment
Of the Requirements
For the Degree of

Master of Science

July 2015

I have submitted this thesis in partial fulfillment of the requirements for the degree of Master of Science.

7/7/15
Date

Marisa L. Naciuk
Marisa L. Naciuk

We approve the thesis of Marisa L. Naciuk as presented here.

7/7/15
Date

Kevin S. Burgess
Kevin S. Burgess, Associate Professor
of Biology & Thesis Advisor,
Columbus State University

7/3/15
Date

Laura F. Galloway
Laura F. Galloway, Professor of Biology,
University of Virginia

1 July 2015
Date

Holly R. Prendeville
Holly R. Prendeville, Research
Geneticist, USDA Forest Service, OR

TABLE OF CONTENTS

| | |
|--|------|
| Table of Contents..... | iii |
| List of Tables | iv-v |
| List of Figures..... | vi |
| Acknowledgements..... | vii |
| Abstract..... | 1 |
| Introduction..... | 2 |
| Methods..... | 6 |
| Study System | 6 |
| Common Garden Design..... | 6 |
| Creating experimental seeds | 6 |
| Seed germination experiment | 7 |
| Rosette transplant experiment..... | 9 |
| Measurements of Phenological and reproductive traits | 10 |
| Statistical Analyzes..... | 11 |
| Assessing local adaptation | 11 |
| Phenotypic selection | 12 |
| Hybrid Breakdown..... | 13 |
| Results..... | 13 |
| Discussion..... | 18 |
| Literature Cited | 24 |
| Appendix A..... | 48 |
| Appendix B1 | 49 |
| Appendix B2 | 50 |
| Appendix C | 51 |
| Appendix D..... | 52 |
| Appendix E | 53 |
| Appendix F..... | 54 |
| Appendix G..... | 55 |

LIST OF TABLES

| Table | Page |
|---|------|
| <p>1 Nested ANOVA of minimum number of days to germinant emergence and percent germination across nine parental <i>C. americanum</i> populations, five from a northern origin and four from a southern origin, planted into common gardens at sites in Georgia (south) and Michigan (north). The experiment was repeated in two years. F-values are listed for each fixed and random effects. Block(Site, Year), Population(Origin) are the random effects in this model. There are seven degrees of freedom associated with Population[Origin] and 104-111 degrees of freedom for Block(Site, Year). Degrees of freedom are one for each fixed effect. Denominator d.f. are in brackets. (*)P<0.10, *P<0.05, **P<0.01, ***P<0.001.....</p> | 35 |
| <p>2 ANOVA of reproductive and phenological traits of northern and southern populations of <i>C. americanum</i> planted at Georgia and Michigan common garden sites (2014). F-values are listed for each fixed and random effects. Block[Site] and Pop[Origin] are the random effects in this model. There are seven degrees of freedom associated with Pop[Origin] and 36 degrees of freedom for Block[Site]. Degrees of freedom are one for each fixed effect (*)P<0.10, *P<0.05, **P<0.01, ***P<0.001.....</p> | 36 |
| <p>3 ANOVA of cumulative fitness of northern and southern origins planted at Georgia and Michigan common garden sites (2014). F-values are listed for each fixed and random effects. Block[Site] and Pop[Origin] are the random effects in this model. There are seven degrees of freedom associated with Pop[Origin] and 38 degrees of freedom for Block[Site]. Degrees of freedom are one for each fixed effect.....</p> | 37 |

LIST OF TABLES

| Table | Page |
|-------|---|
| 4 | Standardized selection differentials (S), linear gradients (β), and quadratic gradients (γ) of northern and southern <i>Campanulastrum americanum</i> populations planted in common garden sites located in Columbus, GA and Hastings, MI. To detect differences in selection across common garden sites, site and gradient interactions were analyzed. Population was included as a blocking term. The F values are reported for these ANCOVAs. Site means: GA (29.58), MI (16.09). (* $P < 0.10$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).....38 |
| 5A | ANOVA testing the effects of generation on expression of reproductive traits of <i>C. americanum</i> populations. The numerator degrees of freedom is one. Each d.f denominator is noted in brackets and F-ratio is listed for each effect. Site, Generation, and Site x Generation are fixed effects. Generation x population is a random effect.39 |
| 5B | Linear Contrast of L.S. means generated by ANOVAs for reproductive traits that quantifies the level of reproductive fitness exhibited by each generation (P, F2, RF2) of <i>C. americanum</i> populations. Reciprocal crosses were also included these analyzes and were noted by placing a capital R in front of the generations associated with the cross. P-values indicate whether F2 (F2 + RF2) means are significantly different from parents. The numerator d.f. is one. An arrow in the mean comparison column indicates if F2s had means lower or higher than the parents.....40 |
| | Computing log ₁₀ 1-transformed cumulative fitness means of source origin within each common garden site. Cumulative fitness was calculated by multiplying total number of fruits by number of seeds per fruit. Data collected from the source common garden experiment (2014) was used in this calculation.48 |
| | Assessing hybrid breakdown by comparing L.S. means (40-45) and number of flowers, (b) total number of fruits, (c) number of seeds/fruit, and (d) biomass. Linear contrast was performed to detect significance between means of parents and F2s of <i>C. americanum</i> populations.47 |

LIST OF FIGURES

| Figure | Page |
|--------|---|
| 1 | Location of nine <i>Campanulastrum americanum</i> populations (AL, NE, MI, TN, OK, MN, MS, WI, and OH) planted in common garden sites, Pierce Cedar Creek Institute in Hastings (MI) and CSU's Oxbow Meadows Environmental Learning Center in Columbus (GA). Common garden site locations are noted with stars. Red lines indicated which populations were crossed to generate F1s and F2s.....41 |
| 2 | Comparing L.S. mean (\pm S.E.) A) minimum number of days to germination, B) percent germination across northern and southern <i>C. americanum</i> populations at Georgia and Michigan common garden sites. Means reflect germination data that was collected for a period of two years (2012-2014).....42 |
| 3 | Assessing variations in phenology. Comparing log+1-transformed mean (\pm S.E.) A) bolting rate, B) number of days to first flower, and C) minimum number of days to fruit maturation across northern and southern <i>C. americanum</i> populations at Georgia and Michigan common garden sites (2014).....43 |
| 4 | Assessing traits indicating size. Comparing log+1-transformed mean (\pm S.E.) A) height at flowering, B) total branch length, and C) biomass across northern and southern <i>C. americanum</i> populations at Georgia and Michigan common garden sites (2014).....44 |
| 5 | Assessing variation in reproductive traits. Comparing log+1-transformed mean (\pm S.E.) A) total number of flowers, B) total number of fruits, C) seeds per fruit, and across northern and southern <i>C. americanum</i> populations at Georgia and Michigan common garden sites (2014)..... 45 |
| 6 | Comparing log+1-transformed cumulative fitness means of source origins within each common garden site. Cumulative fitness was calculated by multiplying total number of fruits by number of seeds per fruit. Data collected from the rosette common garden experiment (2014) was used in this calculation.....46 |
| 7 | Assessing hybrid breakdown by comparing L.S. means (\pm S.E.) A) total number of flowers, B) total number of fruits, C) number of seeds/fruit, and D) biomass. Linear contrast was performed to detect significance between means of parents and F2s of <i>C. americanum</i> populations.....47 |

ACKNOWLEDGEMENTS

It has been a tremendous honor and blessing to work with each of my thesis committee members. They have provided much guidance, support, and encouragement, while navigating through the scientific process. Your wealth of knowledge in the field of ecological genetics has been insightful and your dedication to my success has been greatly appreciated. This opportunity to complete a Masters at CSU has been an important stepping stone towards my career in the field of genetics and will greatly assist in my doctoral studies at University of Georgia. To Dr. Kevin Burgess, my thesis advisor and fellow Canadian, thank you for presenting this opportunity and introducing me to this field of study. I am grateful for your assistance with field work, and guidance with writing this manuscript. To Drs. Laura Galloway and Holly Prendeville, thank you for being so instrumental in the execution of this project, and answering my numerous questions about methodology and data analyzes.

I would also to extend my gratitude to the faculty and students of the Columbus State University Biology Department. Thank you for your encouragement, support, and expressed interest in my research. It has been a pleasure and honor to be part of this department for both my undergraduate and graduate studies. I would especially like to thank these individuals; Samantha Worthy, Scott Silvis, and Lauren Whitehurst. Thank you for spending countless hours in the field, assisting with data collection and garden maintenance. Also, an immense amount of support was given by University of Virginia Biology department. The efforts of the undergraduate students of the Galloway lab need to be recognized. I appreciate your commitment and essential role in data collection and entry for this project.

Financial support was provided by the National Science Foundation, Georgia Native Plant Society, and Columbus State University Research and Creative Endeavors Committee. Thank you for electing to fund my research, allowing essential materials to be purchased. I would also like to thank CSU's Oxbow Meadows Environmental Learning Center for materials and site access. Dr. George Stanton and Oxbow staff thank you for always being very helpful and accommodating. All these organizations have provided essential assistance needed for the successful completion of this thesis research.

Last but not least, to my family and friends, thank you for being my cheerleaders, therapists, and life coaches. Thank you for always believing in my potential, supporting my career aspirations, and never hesitating to help when needed, even if that meant assisting with field work in the rain or intense heat. You have helped me keep grounded during this journey, always reminding me that patience, motivation, and hard work always pays off. I am extremely blessed to have your love and support in my life.

ABSTRACT

Local adaptation is characterized by higher reproductive fitness by individuals present in their native habitats relative to nonlocals. This study utilized *Campanulastrum americanum* as a model for assessing whether an environmental gradient can facilitate local adaptation. In particular, I determined if a latitudinal gradient across eastern United States has influenced the phenology and reproductive fitness of five northern and four southern *C. americanum* populations. These populations were reciprocally transplanted into two common garden sites positioned in the most northern and southern extent of the species distribution: Hastings, MI and Columbus, GA. I observed adaptive population differentiation of northern and southern populations. Both reproductive and phenological traits of plants of southern origin were differentiated from plants of northern origins and under selection. There was also significant selection towards earlier flowering initiation. Earlier flowering, along with faster bolting rate and delayed fruit maturation has assisted in maximizing reproductive fitness of southern populations in the south. While, earlier flowering, delayed fruit maturation, and delayed bolting has facilitated higher fitness of northern populations in the northern part of the species range. This study demonstrates that selection may occur across a species distribution, creating locally adapted populations.

INTRODUCTION

Local adaptation has an influential role in maintenance of genetic and phenotypic diversity, facilitating ecological speciation, and expanding species ranges (Levene 1953; Felsenstein 1976; Hendrick 1986; Kirpatrick & Barton 1997; Tiffin & Ross-Ibarra 2014). It is characterized by a higher level of Darwinian fitness expressed by individuals present in their native habitats relative to foreigners (William 1966). The occurrence of local adaptation is dependent on the interactions between evolutionary forces (ie: selection, gene flow, drift, and mutations), and the presence of heterogeneous environmental conditions that create variation in selection pressures (Blanquart *et al.* 2013). Studies have utilized geographical variations, such as latitudinal clines to address how local adaptation can relate to environmental variation (Mitchell-Old *et al.* 2007; Svetec *et al.* 2015).

Environmental factors can differ throughout a species distribution, creating spatial heterogeneity among populations; thereby exerting selective pressures that can drive divergent changes in population genetic structure among populations (Hunter 2006; Coop *et al.* 2010; Eckert *et al.* 2010; Tanja *et al.* 2013). These changes reflect adaptive gene complexes that confer benefits (higher fitness) for a phenotype in particular set of environmental conditions (Levene 1953; Felsenstein 1976; Hedrick *et al.* 1976; Hedrick 1986; Galloway & Fenster 2000; Kawecki & Ebert 2004; Riis *et al.* 2010; Anderson *et al.* 2011; Luquet *et al.* 2015). Overtime, the frequency of these changes accumulates, forming genetically and physiologically distinct populations (Hunter 2006). Thus, environmental variation has major implications on genetic and subsequent phenotypic differentiation of organisms (Miller & Fowler 1994; Ward *et al.* 2012). Phenotypic

variation due to underlying genetic differences among individuals can serve as evidence of populations adapting to their local environments (Clausen *et al.* 1940; Linhart & Grant 1996; Miaud & Merila 2001; Garcia *et al.* 2007; Feder & Nosil 2010; Blanquart *et al.* 2012; LeCorre & Kremer 2012).

Phenotypic variation can also be a by-product of differences in genotype expression in response to environmental heterogeneity, also referred to as phenotypic plasticity (Scheiner 1993; Scheiner & Lyman 1991; Agrawal 2001; Pigliucci 2005; Gratani 2014). The advantage of plasticity is that multiple phenotypes can be expressed depending on the environment, allowing organisms to deal with unpredictability (Fordyce 2006; Ghalambor *et al.* 2007; Fusco & Minelli 2010; Scheiner 2014). Most importantly, it provides a variety of phenotypes for natural selection to act upon (Lazzaro *et al.* 2008). Phenotypic plasticity becomes adaptive when a phenotype confers higher fitness within a particular environment that has no associated genetic differentiation (Kawecki & Ebert 2004; Kleunen & Fischer 2005; Volis *et al.* 2005; Beldade *et al.* 2011; Ward *et al.* 2011). The environment drives genetic changes in populations that promote adaptation as well as promote variable gene expression that has the potential to become adaptive as well.

Genetic differentiation of populations may also reflect coordination of phenology with their environment to maximize their fitness (Rathcke & Lacey 1985; Reece & Bazzaz 1987; Kozlowski 1992; Levin 2006). Organisms, such as plants have some indication of conditions present in their environment by relying on specific environmental elements as cues such as temperature and amount of light (Karban 2008). Growth and reproduction phenology is in accordance with these cues to ensure that reproduction occurs in the most appropriate set of environmental conditions. In the case

of a plant, flowering, fruiting and seed dispersal are key aspects of reproduction, and a vital determinant of an organism's fitness (Kim *et al.* 2009).

Gene x environment interaction not only facilitates the occurrence of local adaptation (Kawecki & Ebert 2004), but also influences the fitness of hybrids between locally adapted populations. Hybrids may have fitness higher than either parent (hybrid vigor or heterosis) or their fitness can be lower, outbreeding depression (Darwin 1867; Dobzhansky 1951; Lynch 1991; Muller 1942; Arnold *et al.* 1999; Whitlock *et al.* 2000; Truelli *et al.* 2001; Lippman & Zamir 2007; Orr 1995; Willet 2012). In many plant species, the positive effects of heterozygosity on fitness are usually observed during the F1 generation, due to overdominance or masking of recessive deleterious alleles (Waser & Price 1994; Fenster & Galloway 2000; Waser *et al.* 2000; Willi & Van Buskirk 2005). Outbreeding depression is often more evident in later generations due to epistatic incompatibility (Levin 1978; Lynch 1991; Edmands 2007). This form of genetic incompatibility is described as a reproductive failure, but it is formally referred to as hybrid breakdown (Oka *et al.* 2004; Yasumoto & Yahara 2008; Burton *et al.* 2013). Dobzhansky-Muller model offers an explanation of how and why this genetic incompatibility arises within later generations (Orr 1995; Coyne & Orr 2004; Fitzpatrick 2008). When populations are isolated, some mutations are favored by natural selection and increase in frequency. When populations mate, these mutations interact during recombination and contribute to a reduction in offspring fitness (Felsenstein 1974; Muller 1964; Edmands 2007).

The American Bellflower (*Campanulastrum americanum* (L.) Small) is a plant species with a wide distribution and exhibits diverse morphological and phenotypic traits

(Galloway & Etterson 2005; Etterson *et al.* 2007). This makes this species an ideal model for assessing the relationship between geographic (latitude) and phenotypic variation, and if phenotypic variations indicate adaptive population differentiation. Given the potential for adaptation to occur across its range facilitated by differences in environmental conditions, latitudinal variation in phenological and reproductive traits can serve as lines of evidence of local adaptation for this species. Variation in flowering time has been observed within and among populations of *C. americanum* (Burgess *et al.* 2007; Galloway & Burgess 2009). This variation has likely evolved in accordance to cues present in their local environment that reduces the risk of floral damage and incomplete seed development, and increases the availability of potential mates (Anderson *et al.* 2011).

The overall goal of my study is to determine if environmental differences that reside along a latitudinal gradient has facilitated local adaptation of *C. americanum* populations. The four specific objectives of my study are as follows: 1) Assess if populations of *C. americanum* are locally adapted by comparing reproductive traits across common garden sites, 2) Assess if phenological traits vary across sites and if they are adaptive, 3) To determine if variations in reproductive and phenological traits are consistent with adaptive differentiation, and 4) Determine the fitness of hybrids and magnitude of hybrid breakdown.

METHODS

Study System

Campanulastrum americanum (L.) Small (= *Campanula americana* L., Campanulaceae) is an autotetraploid herb that has an expansive distribution across eastern and central North America (Prendeville *et al.* 2013). Populations are typically found in disturbed habitats and occupied by deciduous forest (Galloway *et al.* 2003, Galloway 2005). Populations exhibit diverse morphological and phenotypic traits such as number of days to seed emergence, number of days to flower initiation, number of branches, number of fruits, and plant size (Kalisz & Wardle 2004; Galloway & Etterson 2005; Etterson *et al.* 2007; Burgess *et al.* 2007; Haggerty & Galloway 2011). Flowering phenology and timing of germination are closely associated with one another in this species. Timing of germination influences when rosettes are vernalized which must occur to initiate bolting, and thus also impacts when flowering initiation will occur.

Creating experimental seeds for Parents and Offspring

Seeds were sampled from nine populations of *C. americanum* along a latitudinal gradient in 2008 (methods noted in Prendeville *et al.* 2013). Five populations were located in the northern (MI, MN, NE, OH, and WI) region of the species range, and the remaining four resided in the southern (AL, MS, OK, and TN) region (Figure 1, Appendix A). Seed collected from each population contained 15-20 families, to account for genetic variation that may exist within each population. In 2011, ten seeds from family per were sowed in planting trays containing 3 Metromix: 1 Turface. Trays were placed in a growth chamber at University of Virginia with 25 °C day/ 14°C night, 12-hr

days for four weeks. If multiple seeds germinated then seedlings were thinned to one. Trays were then placed in a cold room set at 5 °C and 12-hr days for 7 weeks. Seedlings were transplanted to containers and move to a greenhouse. Plants were watered on daily basis and exposed to a 16-hr light cycle. Once individuals developed three flowers or one week had past since the day of first flower, hand pollination occurred. Within a population individuals were crossed between different families to establish a line of parents. A F1 hybrid generation was established by crossing individuals belonging to different populations noted by state abbreviation: WIxMS, MNxOK, NExAL, OHxOK, TNxMN, and MSxMI (Figure 1). Then reciprocal crosses of these F1 population cross-types were also performed using the same set of pollen donors and pollen recipients to generate F2 hybrids. Generations associated with reciprocal crosses were noted with an "R". Fruits were collected when mature and placed in cool storage until seeds were ready to be used for the seed germination experiment in July 2012. A subset of these seeds was also planted in plug trays, following the above procedure in December 2012. After germination seedlings were exposed to 5 °C for 47 days and then utilized in the rosette transplant experiment.

Seed germination experiment

To address questions of whether populations exhibit higher fitness within their native range and if phenological patterns are indicative of adaptive differentiation, a reciprocal transplant experiment was performed. Seeds from each population were planted on 4-5 & 10-11 August 2012, in two common garden sites, respectively: the Pierce Cedar Creek Institute in Hastings, Michigan and Columbus State University's

Oxbow Meadows Environmental Learning Center in Columbus, Georgia. Each site contained 30 blocks (26.7 cm by 34.3cm) consisting of nine sleeves. There is some variation in the number of seeds per replicate amongst the populations. In one set of populations (AL, MI, MS, TN, and WI), 10 seeds were planted in each replicate and represented once in each block. A total of 30 replicates in each site contained a representative of this set of parents ($10 \text{ seeds/replicate} \times 1 \text{ replicate/block} \times 30 \text{ blocks} = 600 \text{ seeds/population/site}$). The remaining population set (NE, OK, MN, and OH) had five seeds per sleeve and one replicate per block ($5 \text{ seeds/replicate} \times 1 \text{ replicate/block} \times 30 \text{ blocks} = 150 \text{ seeds/population/site}$). The total number of germinants present in each replicate and number of days to germination was monitored in Georgia and Michigan on a bi-weekly basis until 31 May & 2 June 2013, respectively. Another round of planting occurred on 21 August 2013 at Pierce Cedar Creek Institute and 24 August 2013 at Oxbow Meadows, and germination was monitored until 25 April 2014 at Oxbow Meadows and 28 May 2014 at Pierce Cedar Creek Institute. During the second year, blocks consisted of five peat pot trays (nine pots per block). Seeds were planted and observed over a similar period as the previous year's experiment.

Two response variables were scored. Percent germination was calculated by dividing the maximum number of germinants that emerged in a germination replicate by the total number of seeds planted. Minimum time to germination was the number of days between when the seeds were planted and when the germinant(s) were first observed.

Rosette transplant experiment

I also transplanted rosettes to assess phenology and reproductive traits independent of any effects of local adaptation on germination. A total of 317 rosettes were transplanted to each common garden site on 3-4 March 2014 at Oxbow Meadows and 11-13 April 2014 at Pierce Cedar Creek Institute. They were randomly assigned to one of eight blocks and planted 25 cm apart. There is an unbalanced design due to uneven germination and survival. AL, MI, MS, TN, and WI had 17-24 replicates per site, populations NE, OK, MN, and OH had 14-16 replicates/site, and MN had 8 replicates/site. Additionally, there were 11-15 replicates for the F2 associated with each parental cross. Survival was monitored weekly until bolting was evident at which point plants were checked daily for initiation of flowering.

Traits were measured to describe reproductive phenology. Height prior to flowering was determined by conducting a census of plant height prior to the opening of the very first flower of the season (19 June 2014 at Oxbow Meadows and 16 July 2014 at Pierce Cedar Creek Institute). A plant was measured from the base to the tip of the meristem. Height at first flower was taken on the day that the first flower on an individual plant was observed as just opened. The ratio of height prior to flowering and height at flowering is an index of bolting rate. Once the very first flower of the season opened, day of first flower was scored every three days. Number of days to first flower was difference between when rosettes were transplanted and when the first flower was observed on an individual. To monitor fruit maturation, a segment of the mainstem was marked off with twist ties on day of first flower. The first open flower was the first node of this segment, and the next four nodes above this node made up the segment. Fruit maturation was

monitored on a weekly basis in this marked segment until 31 October 2014 at Oxbow Meadows and 2 November 2014 at Pierce Cedar Creek Institute. Minimum number of days to fruit maturation was the difference between when a flower first appeared and when maturation was first noted in the marked node region.

I also measured a suite of size and reproductive traits. Total branch length was calculated by adding the length of all the branches on an individual plant measured on the day of first flower. Flower production was monitored on a weekly basis. The total number of flowers was calculated by summing the total number of opened flowers scored each week until 31 October 2014 at Oxbow Meadows and 2 November 2014 at Pierce Cedar Creek Institute. Two near-mature fruits were collected, i.e. brown in color but pores not open. Fruits were collected at random from one or two nodes above or below the marked node region. The total numbers of seeds per fruit was determined by taking the average of seeds produced by two fruit samples.

Plants were harvested when 80-100% of fruits in the marked node region were mature. Plants were cut at the very base of the stem, just above the ground. If conditions were wet, the plants were placed in a drier (85°F) for 3-5 days. On the other hand, if conditions were very dry, drying was not required. Biomass was determined by weighing dried harvested plants. Fruit number was determined by counting fruits on harvested plants as well as any fruits that fallen off and present at the bottom of collection bags.

If a weekly census was missed, it was conducted a day after the designated census date. If a three day census was missed it was conducted on the next scheduled date, and it was noted that data reflected flowering activity that had occurred over two census

periods. Lastly, any measurements of individual plants that were not taken on the day of first flower were collected on the next scheduled three day census.

Statistical Analyzes

Assessing local adaptation

Phenological, size and fitness traits were compared across common garden sites and populations using an ANCOVA (JMP Statistical software Version 12, SPSS Statistics Version 23). The model included fixed effects of “origin,” whether populations originated from the north or south, “site” as planting location, and random effects population nested in origin and block nested in site. For germination traits, “year” was also included as a fixed effect and block was nested in year and site. An origin x site interaction would provide evidence that performance of populations of each origin differed between sites. Local adaptation would be supported if populations native to the region exhibit higher reproductive fitness than populations from different latitudes. Tukey HSD was used to assess which traits means were statistically different within and across sites. To meet assumptions of normality, days to germination and percent germination were log-transformed. All remaining reproductive and phenological traits were log+1-transformed.

I also assessed local adaptation by calculating cumulative fitness of individuals from different source origins by multiplying the number of seeds per fruit by total number of fruits. An ANOVA was conducted to compare cumulative fitness associated with each origin within and across sites. Site, origin, and site x origin were fixed effects, while population nested in origin and block nested in site were random effects.

Phenotypic selection analysis

A phenotypic selection analysis was performed to assess if phenotypic plasticity observed in each common garden site was adaptive (Lande & Arnold 1983), and if patterns of selection varied across sites. Nine parental populations were included in this analysis. Both phenological (number of days to first flower and bolting rate), size (biomass, total branch length, and height at flowering), and reproductive (total number of flowers, seeds/fruit) traits were included. All of these traits were first standardized to a mean of zero and unity of variance prior to conducting the analysis. I also assessed for correlations and multicollinearity among the traits. Pearson product-moment correlation coefficients revealed that total number of fruits was significantly correlated with biomass, total number of flowers, and height at flower (Appendix B1, B2). Variance inflation factors generated were all less than three, indicating that results of this analysis was not influenced by multicollinearity that exists among these traits

Total number of fruits was used as measure of relative fitness. Fruit counts at each site were divided by the associated site mean. Standardized linear selection gradients (β), were generated by performing a multiple regression of relative fitness on standardized traits to obtain partial regression coefficients. Standardized nonlinear selection gradients (γ) are doubled parameter estimates generated from a multiple regression of relative fitness on standardized traits and their squares. Population was included as blocking term for both multiple regression analyzes. To test whether each gradient (selection) significantly varied across sites ANCOVAs were performed, in which standardized phenological traits were assigned as covariates and common garden site was a fixed effect.

Assessment of hybrid breakdown

To test whether hybrid breakdown occurred in between population crosses, an ANOVA (JMP Statistical software Version 12.0) was performed with generation, cross, site, generation x site, generation x cross, and cross x site as fixed effects. A linear contrast was performed on the L.S. means of each level of generation associated with a particular trait. Fitness traits total number of flowers, total number of fruits, seeds/fruit, and biomass were included in this assessment. I am most interested in testing whether parents (P1 & P2) in an associated cross outperformed (higher fitness) their hybrid offspring (F2s). Reciprocal crosses were also included and were noted by placing a capital R in front of the generations associated with the cross. A linear contrast: $(P1+P2)/2=(F2+rF2)/2$ was used to test whether the performance of F2 is lower than the average performance of parents.

RESULTS

Germination

Percentage of seedling emergence and time to germination significantly varied between northern and southern populations in the Michigan site but not the Georgia site. Timing of seedling emergence of northern populations was significantly delayed at the Michigan common garden site (mean=160 days), taking an additional 83 days to germinate (Table 1, Figure 2a). Plants of northern origin also took an additional 44 days to germinate than southern populations at the Michigan site (Table 1). Southern populations on average took 66 days to germinate at both sites (Figure 2a). A greater percentage of seedlings emerged from populations of southern origin (mean~13%)

compared to northern origin (mean~10%; Table 2, Figure 2b). The Michigan site exhibited a significantly higher percentage of seedling emergences (mean ~15%) relative to Georgia (mean~7%). At the Michigan site, approximately 5% more southern seedlings emerged compared to seedlings of northern origin (Figure 2b).

Phenology

All phenological traits exhibited a significant site effect and site x origin interaction (Table 1 & 2). Phenology of plants of southern and northern origin varied significantly across sites, suggesting plasticity. Within sites, not all phenological traits exhibited significant variation, for example bolting rate. A significant difference in bolting rate was only observed in Georgia, where northern plants bolted significantly slower than southern plants (N mean~0.21, S mean~0.36). In Michigan, bolting rates of plants from different origins were very similar (N mean~0.44, S mean~0.47; Table 2, Figure 3a). Across sites, populations of northern origin were on average only 0.30 of their height at flowering relative to the 0.41 for southern plants, indicating slower bolting in northern plants.

Unlike bolting phenology, distinct trends were observed in flower initiation and fruit maturation within both sites. Significantly earlier flowering and delayed fruit maturation was exhibited by populations present in their native range (Table 2, Figure 3b,c). Native populations flowered six days sooner. There was variation in how delayed fruit maturation occurred in populations native to each site. Fruits of populations native to the southern common garden site matured 17 days later, while fruits of populations native to the northern site took an additional seven days to mature. Furthermore, there was a notable difference in timing of flowering of plants of southern origin across sites.

Southern plants took longer to flower in Michigan by approximately 10 days, while flowering of plants of northern origin varied by four days across sites (Figure 3b). Additionally, as observed with flowering time, fruit maturation of southern plants varied significantly across sites (Table 2). Fruits of plants of this origin took additional 22 days to mature, while fruit maturation of northern plants varied by four days.

Reproductive Traits

Size traits revealed greater variation between populations of different origin in the Georgia site. Size of plants were greater in Georgia (mean biomass=2.07g, mean height=73cm, mean branch length=35cm) compared to Michigan (mean biomass=0.53g, mean height=54cm, mean branch length=0.02cm). Furthermore, height at flowering and total branch length displayed a significant site x origin interaction (Table 2). Plants of southern origin were 0.24g heavier and 14 cm taller at flowering than northern plants at the Georgia site, and total branch length of plants of this origin was 34.98cm longer (Table 2, Figure 4). In Michigan, size of plants from different source origins was comparable.

Reproductive traits revealed significantly greater reproductive output of populations native to common garden locations. All reproductive traits significantly varied across sites. Approximately 3x more flowers, 3x more seeds, and 6x more fruits were produced by plants from both source origins in Georgia than Michigan (Figure 5). Within sites, native populations produced approximately double the number of seeds and flowers compared to nonlocals. Fruit production of native populations did not significant differ from foreign populations (Figure 5c).

Total fruit number and seeds per fruit were used to estimate cumulative fitness of each source origin. Cumulative fitness of each origin significantly varied across and within sites (Table 3, Figure 6). Populations native to the region, where the common garden resided, had a cumulative fitness that was 1.3x higher than the nonlocals; same for both sites. The fitness levels found in Georgia were 9x higher than the levels found in Michigan.

Phenotypic selection analyzes

There was stronger direct selection (β) on the phenological and reproductive traits of *C. americanum* populations in the Georgia common garden site (Table 4). Earlier flowering time was favored at the Georgia site but no selection on flowering time was found in the Michigan site. There was significant selection towards a larger number of flowers, and biomass at the Georgia site but not in Michigan. Michigan displayed selection for greater height at flowering and smaller seed number, whereas no significant selection on these traits was evident in Georgia. Both Georgia and Michigan displayed significant selection on biomass, though the selection for biomass in Georgia was 1.3x greater (Table 4).

Standardized nonlinear selection was evident in both sites (Table 4). A positive quadratic selection gradient is present for the number of flower produced, total branch length, and biomass at the Georgia site. This suggests that relative fitness (fruit production) is an increasing function of greater flower production and larger plant size (branch length and biomass). There is also negative quadratic selection gradient associated with number of days to first flower (Table 4). Individuals in Georgia that

flower earlier and yield a larger reproductive output will have higher fitness. In the Michigan common garden site, there are also positive quadratic selection gradients evident. Relative fitness is a function of greater height at first flower and biomass, and thus larger size would be associated with higher fitness. The number of seeds per fruit is the only trait that has a negative quadratic selection gradient at the Michigan site, indicating a decelerating relationship between relative fitness and seed number (Table 4). The phenotypic distribution of each trait was reviewed by analyzing graphs. Number of days to first flower and seeds per fruit are under stabilizing selection, indicated by a higher frequency of individuals expressing phenotypes in the middle of the distribution. Height at first flower, total number of flowers, total branch length, and biomass are under disruptive selection, evident by greatest number of individuals expressing a phenotype towards one end of the distribution.

Hybrid breakdown

Biomass indicated that there was a significant generational effect, while the other traits did not. (Table 5A, Figure 7). Linear contrast revealed that there was significant variation in number of seeds per fruit for each level of generation, which indicates hybrid breakdown. Furthermore, there is significant variation in biomass for each generation level that indicates heterosis (Table 5B). When analyzing variation across levels of generation by site and population cross-type, mean differences were consistently insignificant for majority of the traits (Appendix C, D).

DISCUSSION

Local adaptation

Reproductive fitness clearly suggests both northern and southern populations are locally adapted. In this study, fitness is indicated by flower, seed, and fruit production, along side traits tied to size (height at first flower, biomass, and total branch length).

Fitness of native plants was significantly higher at both sites, and thus they are outperforming individuals that are not native to their range. Furthermore, differences in plant size may also be related to reproductive success in a particular environment

(Primack 1987; Wesselingh et al. 1997). Greater size was observed by both origins at the Georgia site, implying that size of plant is more indicative of climatic differences.

Overall larger plant size in the southern common garden, along with heritable phenotypic variation selected by natural selection within southern plants, has resulted in higher fecundity and higher reproductive output. Estimated cumulative fitness values serve as further evidence of local adaptation revealed by reproductive traits, illustrating higher fitness levels by natives as well.

Percent germination was also used an indicator of fitness. Plants of southern origin had the highest percentage germination across sites, displaying higher fitness than northern populations at both sites. Higher fitness of southern plants at the Georgia site and lack of evident of higher fitness of northern plants at the Michigan site is inconsistent with the other reproductive traits.

Underlying differentiation in phenology may facilitate local adaptation. In my study, phenological traits varied significantly across sites for plants of southern and northern origins. This suggests that phenological traits for both source origins are plastic.

Earlier flowering and delayed fruit maturation were exhibited by both origins within their native range. On the other hand, variation in bolting rate was only observed in the Georgia site, in which native populations bolted significantly earlier. Similar to my study, delayed fruit maturation and delayed bolting have been reported in *C. americanum* populations native to regions of northern latitude and higher elevations (Haggerty & Galloway 2011; Prendeville *et al.* 2013). Other plant species such as *Arabidopsis thaliana* demonstrate a positive correlation between number of days to bolt and latitude, where plants from northern latitudes bolted later than plants from southern latitudes.

* Temperature fluctuates in a systematic fashion along a latitudinal gradient (Johanson *et al.* 2000; Stinchcombe *et al.* 2004). Thus, cooler temperatures may support later bolting in this species. Whether plasticity of phenological traits of southern and northern populations is adaptive can be addressed by relating phenology to reproductive fitness. Fitness is higher for populations present in their native range. It can be concluded that earlier flower, earlier bolting, and delayed fruit maturation may have been selected for in southern extent of the distribution, because they maximize the reproductive output of southern plants. While, earlier flowering, delayed fruit maturation, and delayed bolting may assist with increasing the fitness of northern plants.

The role of time to germination in reproductive fitness is unclear. Earlier germination in the Georgia common garden site suggests that southern plants would complete their reproductive cycle faster. Timing of germination may be a reflection of a suite of environmental factors that could influence seedling emergence; light, soil moisture, salinity, and temperature (Tanveer *et al.* 2012). Temperature has been shown to significantly increase phenological progression in warmer temperatures (Gordo & Sanz

2010; Hulber *et al.* 2010; Galloway and Burgess 2009; Haggerty & Galloway 2011), and thus could promote shorter germination time at southern latitudes.

Phenotypic selection

There is selection towards greater number of flowers, biomass, and length of branches in the southern common garden site. While, only a few reproductive traits (height at flower, biomass, and seeds) are under significant selection at the northern site.

This shift towards higher reproductive output (flowers) and overall size (biomass, branches) in the southern site is also observed, indicating that plasticity is adaptive.

Unlike my study, previous experimental research on *C. americanum* have report no relation between biomass and latitude gradient. However, what studies have found that are consistent with my results is that there are significant variations in branch lengths and final plant size (Kalisz & Wardle 1994; Prendeville *et al.* 2013). In other species such as *Arabidopsis thaliana*, biomass has been shown to negatively correlate with elevation with smaller plants associated with cooler environments (Montesinos-Navarro *et al.* 2011). Length of growing season, amount of rain, temperature, and characteristics of vegetation cover can influence plant size (Crauford & Wheeler 2009; Rajasekar *et al.* 2013). A combination of environmental factors existing along a latitudinal gradient may be contributing to overall selection for larger size.

Furthermore, I observed significant selection for earlier flowering in the southern populations in the southern site but not the northern site. The length of time to flower initiation can influence how many flowers or fruits an individual can produce during a growing season (Ollerton & Lack 1998). Earlier flowering may be beneficial in that there

is a longer flowering duration, higher likelihood of fertilization and visitation of pollinators, and reduction in competition with other flowering species; contributing to reproductive success (Mosquin 1971; Rathcke & Lacey 1985; Gentry 1974; Heinrich 1975; Stiles 1975; Alonso 2004). Longer growing seasons present in the south would allow an individual plant to produce more flowers (Haggerty & Galloway 2011). Greater flower production in the south could possibly be facilitated by a longer growing season and earlier flower initiation. This is in accordance with the expectation that longer growing season and warmer temperatures would encourage earlier flower initiation (Fitter & Fitter 2002; Etterson 2004; Griffith & Watson 2005; Parmesan 2007; Gordo & Sanz 2009; Galloway & Burgess 2012). This behavior has also been observed in *C. americanum* at lower elevations, warmer conditions (Haggerty & Galloway 2011). Delayed flowering initiation has also been observed by other species native to southern latitudes such as *Lythium salicaria* (Olsson & Agren 2002).

It has been demonstrated that flowering time can be a plastic, responding to environmental cues such as photoperiod, vernalization, and resource availability that indicate when conditions are best to reproduce (Levy & Dean 1998; Gordo & Sanz 2009). However, flowering time may also be reflective of internal cues (Levy & Dean 1998; Burgess *et al.* 2007; Haggerty & Galloway 2011). The results of this study suggest that flowering time is an adaptive response of southern and northern populations. When populations were closer to their home sites displayed quicker flowering initiation but when placed in a location outside of their native range their flowering period was delayed. *C. americanum*, along with *Melandrium* and *Aradidopsis*, exhibit distinct flowering genotypes across a specified geographic range (Lawrence 1963; Westerman

1971; Stincombe *et al.* 2004). Thus, variation in flowering time could potentially be reflected of genotypes selected in a particular environment.

The magnitude of selection and the traits under selection appear to differ across sites. To confirm whether selectional gradients differed across sites, analysis of covariance was performed to assess if selection gradients differed across the common garden sites. There was significant site x gradient interaction for the total number of days to first flower, total number of flowers, seeds per fruit, height at first flower, total branches, and biomass. This indicates that magnitude of selection on traits at each site varies and this is influencing relative fitness of populations. For majority of these reproductive and phenological traits, except for biomass, there was evidence of local adaptation suggested by higher fitness of populations native to each site and significant site x origin interaction. All of these traits also exhibited a significant site effect. This confirms that the site interaction being observed is due to differences in magnitude of selectional gradients that exist at each site.

Hybrid Breakdown

There is little evidence to suggest that genetic differentiation between northern and southern populations has resulted in hybrid breakdown. Across population cross-types and sites, F2 generations did yield reproductive outputs as large as their associated parents, suggested by linear contrast. This also holds true within each site (Appendix C, D). The results of this study are in accordance with the common observation that few postzygotic barriers to hybridization develop polyploidy. In polyploids, there is a genetic buffer against allele fixation. Any mutations that form in a population are not likely to

become fixed in a population (Stebbins 1950). *C. americanum* is an autotetraploid and if genetic differentiation occurred among populations, a genetic buffer would prevent new mutations from becoming fixed. This could either slow down the formation of a reproductive isolation barrier or limited the possibility of epistatic incompatibility.

Adaptive population differentiation

Adaptive population differentiation occurs when natural selection acts upon heritable phenotypic variation (Linhart & Grant 1996; Miao & Merila 2001; Garcia *et al.* 2007). Selective forces have acted upon reproductive and phenological variations present within *C. americanum* populations of different origins, resulting in genetic and phenotypic differentiation. Varying environmental conditions present along a latitudinal gradient has facilitated differentiation. Furthermore, magnitude of selection was not uniform across the gradient, creating differences in selection across sites. Higher fitness levels at home sites suggest that differentiation between these source origins is adaptive. This study demonstrates that utilizing geographical variations such as latitude can assist in investigating whether locally adapted populations have formed across a species range.

LITERATURE CITED

- Alonso C. 2004. Early Blooming's Challenges: Extended Flowering Season, Diverse Pollinator Assemblage and the Reproductive Success of Gynodioecious *Daphne laureola*. *Ann Bot.* 93: 61-66.
- Anderson JT, Willia JH, Mitchell-Olds T. 2011. Evolutionary genetics of plant adaptation. *Trends Genet* 27: 258-266.
- Agrawal AA. 2001. Phenotypic plasticity in the interaction and evolution of species. *Science* 294: 321-325.
- Arnold ML, Bulger MR, Burke JM. 1999. Natural hybridization: how low can you go and still be important? *Ecology* 80: 371-381.
- Atkins OK, Loveys BR, Atkinson LJ, Pons TL. 2005. Phenotypic plasticity and growth temperature: understanding interspecific variability. *Journal of Experimental Botany* 57: 267-281.
- Beaumont MA. 2005. Adaptation and speciation what can F_{ST} tell us? *Trends in Ecology & Evolution* 20: 435-440.
- Blanquart F, Gandon S, Nuismer SL. 2012. The effects of migration and drift on local adaptation to a heterogeneous environment. *Journal of Evolutionary Biology* 25: 1351-1363.
- Blanquart F, Kaltz O, Nuismer SL, Gandon S. 2013. A practical guide to measuring local adaptation. *Ecology Letters* 16: 1195-1205.
- Bolmgren K, Lonnberg K. 2005. Herbarium data reveal an association between fleshy fruit type and earlier flowering. *Int J Plant Sci* 166: 663-670.
- Burgess KS, Etterson JR, Galloway LF. 2007. Artificial selection shifts flowering phenology and other correlated traits in an autotetraploid herb. *Heredity* 99: 641-648.
- Burton RS, Pereira RJ, Barreto FS. 2013. Cytonuclear genomic interactions and hybrid breakdown. *Annual Review of Ecology, Evolution, and Systematics* 44: 281-302.
- Calcedo AL, Stinchcombe JR, Olsen KW, Schmitt J, Purugganan MD. 2004. Epistatic interaction between *Arbidopsis* FRI and FLC flowering time genes generates a latitudinal cline in a life history trait. *PNAS* 101: 15670-15675.
- Clausen J, Keck DD, Hiesey WM. 1940. Experimental Studies on the Nature of Species. Effect of varied environments on western North American plants. Carnegie Institute of Washington Publications 1940: 1-452.

- Coop G, Witonsky D, Di Rienzo A, Pritchard JK. 2010. Using Environmental Correlations to Identify Loci Underlying Local Adaptation. *Genetics* 185: 1411-1425.
- Coyne J, Orr HA. 2004. *Speciation*. Sunderland (MA): Sinauer Associates.
- Craufurd PQ, Wheeler TR. 2009. Climate change and the flowering time of annual crops. *Journal of Experimental Botany* 60: 2529-2539.
- Darwin C. 1867. *The effects of cross and self fertilization in the vegetable kingdom*. London: J. Murray.
- Despland E, Houle G. 1997. Climate influences on growth and reproduction of *Pinus banksiana* (Pinaceae) at the limit of the species distribution in eastern North America. *American Journal of Botany* 84: 928-937.
- Dijk HV. 2009. Evolutionary change in flowering phenology in the iteroparous herb *Beta vulgaris* ssp. *Maritima* a search for the underlying mechanisms. *J. Exp. Bot* 60: 3143-3155.
- Dobzhansky T. 1936. Studies on hybrid sterility. II. Localization of sterility factors in *Drosophila pseudoobscura* hybrids. *Genetics* 21: 113-135.
- Dobzhansky T. 1951. *Genetics and the Origin of Species*. 3ED. New York: Columbia University Press.
- Dough Y, Cheng Z, Meng H, Liu H, Wu C, Rehman A. 2013. The effect of cultivar, sowing date and transplantation in field on bolting of Welsh onion (*Allium fistulosum* L.). *BMC Plant Biology* 13: 1-12.
- Edmands S. 2007. Between a rock and a hard place: evaluating the relative risks of inbreeding and outbreeding for conservation and management. *Molecular Ecology* 16: 463-475.
- Etterson JR, Keller SR, Galloway LF. 2007. Epistatic and cytonuclear interaction govern hybrid breakdown in the autotetraploid *Campanulastrum americanum*. *Evolution* 61: 2671-2683.
- Evanhoe L, Galloway LF. 2002. Floral Longevity in *Campanula americana* (Campanulaceae): A comparison of morphological and functional gender phases.
- Felsenstein. 1974. The evolutionary advantage of recombination. *Genetics* 78: 737-756.
- Felsenstein 1976. The theoretical population genetics of variable selection and migration. *Annu. Rev. Genet.* 10: 253-280.

- Fenner M. & Thompson K. 2005. The Ecology of Seeds, 2nd edn, Cambridge University Press, Cambridge, UK
- Galloway 2005. Maternal effects provide phenotypic adaptation to local environmental conditions. *New Phytologist* 166: 93-100.
- Galloway LF, Cirigliana T, Gremski K. 2002. The contribution of display size and dichogamy to potential geitonogamy in *Campanula americana*. *Int. J. Plant Sci* 163: 133-139.
- Galloway LF, Burgess KS. 2012. Artificial selection on flowering time: influence on reproductive phenology across natural light environments. *Journal of Ecology* 100: 852-861.
- Galloway LF, Burgess KS. 2009. Manipulation of flowering time: phenological integration and maternal effects. *Ecology* 90: 2139-2148.
- Galloway LF, Etterson JR. 2005. Population differentiation and hybrid success in *Campanula americana*: geography and genome size. *Journal of Evolutionary Biology* 18: 81-89.
- Galloway LF, Etterson JR, Hamrick JL 2003. Outcrossing rates and inbreeding depression in the herbaceous autotetraploid *Campanula americana*. *Heredity* 90: 308-315.
- Galloway LF, Fenster CB. 1999. The Effect of Nuclear And Cytoplasmic Genes On Fitness And Local Adaptation In An Annual Legume, *Chamaecrista fasciculata*. *Evolution* 53: 1734-1743.
- Galloway LF, Fenster CB. 2000. Population differentiation in an annual legume: local adaptation. *Evolution* 54: 1173-1181.
- Garcia de Leaniz C, Fleming IA, Einum S, Verspoor E, Jordan WC, Consuegra S, Aubin-Horth N, Lajus D, Letcher BH, Youngson AF, Webb JH, Vollestad LA, Villanueva B, Ferguson A, Quinn TP. 2007. A critical review of adaptive genetic variation in Atlantic salmon: implications for conservation. *Bio Rev* 82: 173-211.
- Ghalambor CK, McKay JK, Carroll SP, Reznick DN. 2007. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology* 21: 394-407.
- Galloway LF, Etterson JR, McGlothlin JW. 2009. Contribution of direct and maternal genetic effects to life-history evolution. *New Phytologist* 183: 826-838.
- Gratani L. 2014. Plant Phenotypic Plasticity in Response to Environmental Factors. *Advances in Botany* 208747: 1-17.

- Eckert A, Bower AD, Gonzalez-Martinez SC, Wegrzyn JL, Coop G, Neale DB. 2010. Back to nature: ecological genomics of loblolly pine (*Pinus taeda*, Pinaceae). *Molecular Ecology* 19: 3789-3805.
- Engelen-Eigles G, Erwin JE. 1997. A model plant for vernalization studies. *Sci Hortic* 70: 197-202.
- Etterson JR. 2004. Evolutionary potential of *Chamaecrista fasciculata* in relation to climate change. Clinal patterns of selection along an environmental gradient in the Great Plains. *Evolution* 58: 1446-1458.
- Feder JL, Nosil P. 2010. The efficacy of divergence hitchhiking in generating genomic islands during ecological speciation. *Evolution* 64:1729-1747.
- Felsenstein J. 1976. The theoretical population genetics of variable selection and migration. *Annu. Rev. Genetics* 10: 253-280.
- Fitter AH, Fitter RSR. 2002. Rapid changes in flowering time in British plants. *Science* 296: 1689-1691.
- Fordyce, JA. 2006. The evolutionary consequences of ecological interactions mediated through phenotypic plasticity. *J. Exp. Biol.* 209: 2377-2383.
- Fusco G, Minelli A. 2010. Phenotypic plasticity in development and evolution: facts and concepts. *Philos Trans R Soc Lond B Biol Sci.* 365: 547-556.
- Garcia D, Zamora R, Gomez JM, Jordano P, Hodar JA. 2000. Geographical variation in seed production, predation and abortion in *Juniperus communis* throughout its range in Europe. *Journal of Ecology* 88: 436-446.
- Gentry AH. 1976. Bignoniaceae of southern Central America: distribution and ecological specificity. *Biotropica* 8: 117-131.
- Gomez-Mena C, Pineiro M, Franco-Zorrilla JM, Salinas J, Coupland G, Martinez-Zapater JM. 2001. early bolting in short days: An *Arbidopsis* Mutation That Causes Early Flowering and Partially Suppresses the Floral Phenotype of leaf. *The Plant Cell* 13: 1011-1024.
- Gordo O, Sanz JJ. 2009. Long-term temporal changes of plant phenology in the Western Mediterranean. *Global Change* 15: 1930-1948.
- Gordo O, Sanz JJ. 2010. Impact of climate change on plant phenology in Mediterranean ecosystem. *Global Change Biology* 16: 1082-1106.

Graeber K, Nakabayashi K, Miatton E, Leubner-Metzger G, Soppe WJ. 2012. Molecular mechanisms of seed dormancy Plant Cell Environ 35: 1769-1786.

Griffith TM, Watson MA. 2005. Stress avoidance in a common annual: reproductive timing is important for local adaptation and geographic distribution. Journal of Evolutionary Biology 18: 1601-1612.

Felsenstein J 1979. Excursions along the interface between disruptive and stabilizing selection. Genetics 93: 773-795.

Fitzpatrick BM. 2008. Dobzhansky-Muller model of hybrid dysfunction supported by poor burst-speed performance in hybrid tiger salamanders. Journal of Evolutionary Biology 21: 342-351.

Haggerty BP, Galloway LF. 2011. Response of individual components of reproductive phenology to growing season length in a monocarpic herb. Journal of Ecology 99: 242-253.

Hahn MA et al. 2012. Increased Phenotypic Plasticity to Climate May Have Boosted the Invasive Success of Polyploid *Centaurea stoebe*. PLoS ONE 7: e50284.

Hedrick PW. 1986. Genetic polymorphism in heterogeneous environments: a decade later. Annu. Rev. Ecol. Syst. 17: 535-566.

Hedrick PW, Ginevan ME, Ewing EP. 1976. Genetic polymorphism in heterogeneous environments. Ann Rev Ecol. Syst 7: 1-32.

Heinrich B. 1975. Bee flowers: a hypothesis on flower variety and blooming times. Evolution 29: 325-334.

Hulber K, Sonnleitner M, Flatscher R, Berger A, Dobrovsky R, Niessner S, Nigl T, Schneeweiss GM, Kubesova M, Rauchova J, Suda J, Schonswetter P. 2009. Ecological segregation drives fine-scale cytotype distribution of *Senecio carniolicus* in the Eastern Alps. Preslia 2009: 309-319.

Hunter, P. 2006. Defining Species: The indirect impact of humans on biodiversity. EMBO reports 7: 763-766.

Hutchings JA. 2011. Old wine in new bottles: reaction norms in salmonid fishes. Heredity 106: 421-437.

Jong, MD, Leyser O. 2012. Developmental Plasticity in Plants. Cold Spring Harb Symp Quant Biol 77: 63-73.

- Kalisz S, Wardle, GM. 1994. Life history Variation in *Campanula americana* (Campanulaceae): Population Differentiation. *American Journal of Botany* 81: 521-527.
- Karban R. 2008. Plant behavior and communication. *Ecology Letters* 11: 727-739.
- Kilkenny FF, Galloway LF. 2008. Reproductive success in varying light environments: direct and indirect effects of light on plants and pollinators. *Oecologia* 155: 247-255.
- Kim D-H, Doyle MR, Sung S, Amasino RM. 2009. Vernalization: Winter and the Timing of Flowering in Plants. *Annu. Rev. Cell Dev. Biol.* 25: 277-299.
- Kirkpatrick M, Barton N. 1997. Evolution of a species range. *Am. Nat.* 150: 1-23.
- Kawecki TJ, Ebert D. 2004. Review: Conceptual issues in local adaptation. *Ecology Letters* 7: 1225-1241.
- Khokhar KM. 2009. Effect of set-size and storage temp on bolting, bulbing, and seed yield in two onion cultivars. *Sci Horti* 122: 187-194.
- Kozlowski J. 1992. Optimal allocation of resources to growth and reproduction: implications for age and size at maturity. *Trends Ecol. Evol.* 7: 15-18.
- Laporte V, Cugen J, Couvet D. 2000. Effective Population Sizes for Cytoplasmic and Nuclear Genes in a Gynodioecious Species: The Role of the Sex Determination System. *Genetics* 154: 447-458.
- Lawrence CW. 1963. Genetics studies on wild populations of *Melandrium* II. Flowering time and plant weight. *Heredity* 18: 149-164.
- Lazzaro BP, Flores HA, Lorigan JG, Yourth CP. 2008. Genotype-by-Environment Interactions and Adaptation to Local Temperature Affect Immunity and Fecundity in *Drosophila melanogaster*. *PLoS Pathogens* 4: e1000025
- Le Corr V, Kremer A. 2012. The genetic differentiation at quantitative trait loci under local adaptation. *Molecular Ecology* 21: 1548-1566.
- Levin DA. 1978. The origin of isolating mechanisms in flowering plants. *Evolutionary Biology* 11: 185-317.
- Levene H. 1953. Genetic equilibrium when more than one ecological niche is available. *Am. Nat.* 87: 331-333.

- Levene H. 1953. Genetic equilibrium when more than one ecological niche is available. *Am. Nat.* 87: 331-333.
- Levin DA. 2006. Flowering phenology in relation to adaptive radiation. *Syst. Bot.* 31: 239-246.
- Levy YY, Dean C. 1998. The Transition to Flowering. *The Plant Cell* 10: 1973-1989.
- Lippman ZB, Zamir D. 2007. Heterosis: revisiting the magic. *Trends Genet* 23: 60-66.
- Linhart YB, Grant MC. 1996. Evolutionary significance of local genetic differentiation in plants. *Annual Review of Ecology and Systematics* 27: 237-277.
- Luquet E, Lena JP, David, P, Prunier J, Joly P, Lengagne T, Perrin N, Plenet S. 2013. Within-and among-population impact of genetic erosion on adult fitness-related traits in the European tree frog *Hyla arborea*. *Heredity* 110: 347-354.
- Lynch M. 1991. The genetic interpretation of inbreeding depression. *Evolution* 45: 622-629.
- Miaud C, Merila J. 2001. Local adaptation or environmental induction? Cause of population differentiation in alpine amphibians. *Biota* 2: 31-50.
- Michaels SD, Amasino RM. 2000. Memories of winter: vernalization and the competence to flower. *Plant, Cell and Environment* 23: 1145-1153.
- Milford GFJ, Jarvis PJ, Walters C. 2010. A verbalization-intensity model to predict bolting in sugar beet. *J. Agr Sci* 148: 127-137.
- Miller RE, Folwer NL. 1994. Life History Variation and Local Adaptation Within Two Populations of *Bouteloua Rigidiesta* (Texas Grama). *Journal of Ecology* 82: 855-864.
- Mitchell-Olds T, Willis JH, Goldstein DB. 2007. Which evolutionary processes influence natural genetic variation for phenotypic traits? *Nat Rev Genet* 8: 845-856.
- Mosquin T. 1971. Competition for pollinators as a stimulus for evolution of flowering time. *Oikos* 22: 398-402.
- Muller. 1942. Isolating mechanisms, evolution and temperature. *Biological Symposium* 6: 71-125.
- Muller HJ. 1964. The relation of recombination to mutational advance. *Mutat. Res.* 1:2-9.
- Oka, A et al. 2004. Hybrid Breakdown Caused by Substitution of the X Chromosome Between Two Mouse Subspecies. *Genetics* 166: 913-924.

- Ollerton J, Lack A. 1998. Relationships between flowering phenology, plant size and reproductive success in *Lotus corniculatus* (Fabaceae). *Plant Ecology* 139: 35-47.
- Olsson K, Agren J. 2002. Latitudinal population differentiation in phenology, life history and flowering morphology in the perennial herb *Lythium salicaria*. *J. Evol Bio* 15: 983-996.
- Orr HA. 1995. The population genetics of speciation: the evolution of hybrid incompatibilities. *Genetics* 139: 1803-1813.
- Papaix J, David O, Lannon C, Monad I. 2013. Dynamics of adaptation in Spatially Heterogeneous Metapopulations. *PLos ONE* 8: e54697.
- Parmesan C. 2007. Influences of species, latitudes and methodologies on estimates of phenological response to global warming. *Global Change Biology* 13: 1860-1872.
- Penuelas J, Filella I, Comas, P. 2002. Changed plant and animal life cycles from 1952 to 2000 in the Mediterranean region. *Global Change Biology* 8: 531-544.
- Pigliucci M. 2005. Evolution of phenotypic plasticity: where are we going now?. *Trends Ecol Evol* 20: 481-486.
- Post ES, Pederson C, Wilmers CC, Forchhammer MC. 2008. Phenological sequences reveal aggregate life history response to climatic warming. *Ecology* 89: 363-370.
- Primack RB. 1987. Relationships among flowers, fruits, and seeds. *Annual Review of Ecology and Sytematics* 18: 409-430.
- Prendeville HR, Barnard-Kubow K, Dai C, Barringer B, Galloway LF. 2013. Clinal variation for some phenological traits across a species range. *Oecologia* 173: 421-430.
- Rajasekar M, Arumugam T, Ramesh Kumar S. 2013. Influence of weather and growing environment on vegetable growth and yield. *Journal of Horticulture and Forestry* 5: 160-167.
- Rathcke B, Lacey EP. 1985. Phenological patterns of terrestrial plants. *Annual Review of Ecology, Evolution and Systematics* 16: 179-214.
- Reed TE, Schindler DE, Waples RS. 2011. Interacting Effects of Phenotypic Plasticity and Evolution on Population Persistence in a Changing Climate. *Conserv Biol* 25: 56-63.
- Reeekie EG, Bazzaz FA. 1987. Reproductive efforts in plants. *Am. Nat.* 129: 876-919.

- Riis T, Lambertini C, Olesen B, Clayton JS, Brix H, Sorrell BK. 2010. Invasion strategies in clonal aquatic plants: are phenotypic differences caused by phenotypic plasticity or local adaptation. *Annals of Botany* 106: 813-822.
- Scarcelli et al. 2007. Antagonistic pleiotropic effects reduce the potential adaptive value of the *Frigida* Locus. *PNAS* 104: 16986-16991.
- Scheiner SM. 1993. Genetics and evolution of phenotypic plasticity. *Annu. Rev. Ecol. Syst.* 24: 35-68.
- Scheiner SM. 2014. The genetics of phenotypic plasticity. XIII. Interactions with developmental stability. *Ecology and Evolution* 4: 1347-1360.
- Scheiner SM, Lyman RF. 1991. The genetics of phenotypic plasticity. II. Responses to selection. *J. Evol. Biol.* 4: 23-50.
- Schluter D. 1988. Estimating the form of natural selection on a quantitative trait. *Evolution* 42: 849-861.
- Stebbins GL. 1950. *Variation and Evolution in Plants*. Columbia University Press; New York and London.
- Stiles FG. 1975. Ecology, flowering phenology and hummingbird pollination of some Costa Rican *Heliconia* species. *Ecology* 56: 285-301.
- Stinchcombe JR, Cynthia W, Ungerer M, Olsen KM, Mays C, Halldorsdottir SS, Purugganan MD, Schmitt J. 2004. A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the flowering time gene *FRIGIDA*. *PNAS* 101: 4712-4717.
- Stinchcombe JR, Caicedo AL, Hopkins R, Mays C, Boyd EW, Purugganan MD, Schmitt J. 2005. Vernalization sensitivity in *Arabidopsis thaliana* (Brassicaceae): the effects of latitude and *FLC* variation. *Am. J. Bot.* 92: 1701-1707.
- Svetec N, Zhao L, Saelao P, Chiu JC, Begun DJ. 2015. Evidence that natural selection maintains genetic variation for sleep in *Drosophila melanogaster*. *BMC Evolutionary Biology* 15:41.
- Tanja P, Hufford MB, Mezouk S, Ross-Ibarra J. 2013. Complex Patterns of Local Adaptation in *Teosinte*. *Genome Biol Evol* 5:1594-1609.
- Tiffin P, Ross-Ibarra J. 2014. Advances and limits of using population genetics to understand local adaptation. *Cell Systems* 29: 673-680.

- Tsang LM, Chan BKK, Ma KY, Chu KH. 2008. Genetic differentiation, hybridization and adaptive divergence in two subspecies of the acorn barnacle *Tetraclita japonica* in the northwestern Pacific. *Molecular Ecology* 17: 4151-4163.
- Turelli M, Barton NH, JA Coyne. 2001. Theory and speciation. *Trends in Ecology and Evolution* 16: 330-343.
- Valladares F, Matesanz S, Guilhaumon F, Araujo MB, Balaguer L, Benito-Garzon M, Cornwell W, Gianoli E, Mkleunen MV, Naya DE, Nictoria, AB, Poorter H, Zavala MA. 2014. The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecology Letters* 17: 1351-1364.
- Van Kleunen M, Fischer M. 2005. Constraints on the evolution of adaptive phenotypic plasticity in plants. *New Phytologist* 166: 49-60.
- Volis S, Olsvi-Whittaker L, Anikster Y, Mendlinger S. 2005. The influence of space in genetic-environmental relationships when environmental heterogeneity and seed dispersal occurs at similar scale. *American Naturalist* 163: 312-327.
- Yamaski A, Tanaka K, Yoshida M, Miura H. 2000. Effects of day and night temps on flower-bud formation and bolting of Japanese bunching onion (*Allium fistulosum* L.). *J. Jpn Soc Hortic Sci* 69: 40-46.
- Ward D, Shrestha MK, Golan-Goldhirsh A. 2011. Evolution and ecology meet molecular genetics: adaptive phenotypic plasticity in two isolated Negev desert populations of *Acacia raddiana* at either end of a rainfall gradient. *Annals of Botany* 109: 247-255.
- Waser NM, Price MV. 1994. Crossing-distance effects in *Delphinium nelsonii*, outbreeding and inbreeding depression in progeny fitness. *Evolution* 48: 842-852.
- Waser NM, Price MV, Shaw RG. 2000. Outbreeding varies among cohorts of *Ipomopsis aggregate* planted in nature. *Evolution* 54: 485-491.
- Wesselingh RA, Klinkhamer PGL, De Jong TJ, Boorman LA. 1997. Threshold size for flowering in different habitats: Effects of size-dependent growth and survival. *Ecology* 78: 2118-2132.
- Westerman JM. 1971. Genotype-environment interaction and development regulation in *Arabidopsis thaliana* IV. Wild Material; analysis. *Heredity* 26: 383-395.
- Whitlock MC, Ingvarsson PK, Hatfield T. 2000. Local drift load and the heterosis of interconnected populations. *Heredity* 84: 452-457.
- William GC. 1966. *Adaptation and Natural Selection*. Princeton University Press, Princeton.

- Willi Y, Van Buskirk J. 2005. Genomic compatability occurs over a range of parental genetic similarity in an outcrossing plant. *Proc. R. Soc. B Biol. Sci.* 272: 2357-2364.
- Willett, CS. 2012. Hybrid Breakdown Weakens under Thermal Stress in Population Crosses of the Copepod *Tigriopus californicus*. *Journal of Heredity* 103: 103-114.
- Woltereck, R.: 1909, 'Weitere experimentelle Untersuchungen über Artveränderung, speziell über das Wesen quantitativer Artunterschiede bei Daphnien', *Verhandlungen der deutschen zoologischen Gesellschaft* 19: 110-173
- Woodward FL. 1997. Life at the edge: a 14-year study of *Verbena officinalis* population's interactions with climate. *Journal of Ecology* 85: 899-906.
- Yasumoto AA, Yahara T. 2008. Reproductive isolation on interspecific backcross of F1 pollen to parental species, *Hemerocallis fulva* and *H. citrine* (Hemerocallidaceae). *J Plant Res* 121: 287-291.

Table 1. Nested ANOVA of minimum number of days to germinant emergence and percent germination across nine parental *C. americanum* populations, five from a northern origin and four from a southern origin, planted into common gardens at sites in Georgia (south) and Michigan (north). The experiment was repeated in two years. F-values are listed for each fixed and random effects. Block(Site, Year), Population(Origin) are the random effects in this model. There are seven degrees of freedom associated with Population[Origin] and 104-111 degrees of freedom for Block(Site, Year). Degrees of freedom are one for each fixed effect. Denominator d.f. are in brackets. (*)P<0.10, *P<0.05, **P<0.01, ***P<0.001

| Response Variable | Site | Year | Origin | Origin x Site | Origin x Year | Site x Year | Pop (Origin) | Block (Site, Year) |
|---------------------|--------------------|----------------|-------------------|-------------------|----------------|-------------------|------------------|--------------------|
| Days to germination | 28.98*** (381) | 0.06 (385) | 5.08 (391) | 12.36*** (378) | 0.58 (379) | 66.94*** (384) | 6.81*** (271) | 1.57** (271) |
| % germination | 77.06*** (1220) | 0.15 (1214) | 12.93** (1229) | 0.15 (1218) | 0.07 (1214) | 0.02 (1220) | 1.30 (1113) | 1.48** (1113) |

Table 2. ANOVA of reproductive and phenological traits of *C. americanum* populations of northern and southern origin planted at Georgia and Michigan common garden sites (2014). F-values are listed for each fixed and random effects. Block[Site], Pop[Origin] are the random effects in this model. There are seven degrees of freedom associated with Pop[Origin] and 36 degrees of freedom for Block[Site]. Degrees of freedom are one for each fixed effect. Denominator d.f. are in brackets. (*)P<0.10, *P<0.05, **P<0.01, ***P<0.001

| | Trait | Site | Origin | Origin x Site | Pop[Origin] | Block[Site] |
|---------------------------------------|--|---------------------|----------------|-------------------|-------------------|------------------|
| A. Phenological Traits | Bolting rate | 105.33*** (503) | 0.55 (503) | 8.78** (447) | 4.07*** (447) | 1.68** (447) |
| | Number of days to first flower | 11.19** (249) | 0.14 (249) | 20.43*** (204) | 16.89*** (204) | 0.63 (204) |
| | Min number of days to fruit mat | 14.45** (242) | 1.04 (242) | 6.77* (197) | 0.71 (197) | 1.19 (197) |
| | Seeds per fruit | 18.65** (296) | 0.58 (297) | 23.21*** (290) | 5.07*** (252) | 1.04 (252) |
| B. Reproductive Traits | Total number of fruit | 5.21* (272) | 0.02 (273) | 1.17 (228) | 2.43* (228) | 3.68*** (228) |
| | Total number of flowers | 38.26*** (558) | 1.21 (452) | 21.89** (557) | 0.34 (385) | 4.53*** (385) |
| | Height at flowering | 0.03 (249) | 0.90 (249) | 15.19*** (204) | 3.42** (204) | 2.78*** (204) |
| C. Size Traits | Total branch length | 10.84** (356) | 6.07* (356) | 10.95** (311) | 0.97 (311) | 0.59 (311) |
| | Biomass | 50.4967*** (237) | 4.13 (299) | 0.29 (290) | 2.45* (254) | 4.93*** (254) |

Table 3: ANOVA of cumulative fitness of northern and southern origins planted at Georgia and Michigan common garden sites (2014). F-values are listed for each fixed and random effects. Block[Site] and Pop[Origin] are the random effects in this model. There are seven degrees of freedom associated with Pop[Origin] and 38 degrees of freedom for Block[Site]. Degrees of freedom are one for each fixed effect

| Effect | d.f. denominator | F-value | p-value |
|---------------|------------------|---------|---------|
| Site | 369 | 9.52 | 0.0276 |
| Origin | 369 | 0.18 | 0.6813 |
| Site x Origin | 241 | 7.29 | <0.0001 |
| Pop[Origin] | 302 | 4.94 | <0.0001 |
| Block[Site] | 302 | 1.71 | 0.0022 |

Table 4. Standardized selection differentials (S), linear gradients (β), and quadratic gradients (γ) of northern and southern *Campanulastrum americanum* populations planted in common garden sites located in Columbus, GA and Hastings, MI. To detect differences in selection across common garden sites, site and gradient interactions were analyzed. The F values are reported for these ANCOVAs. Site means: GA (29.58), MI (16.09). (*) $P < 0.10$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

| | Trait | B | | | γ | | |
|------------------------|------------------------|-----------|----------|----------------|-----------|----------|-----------------|
| | | GA site | MI site | Site x β | GA site | MI site | Site x γ |
| A. Phenological Traits | Days to first flower | -0.844*** | -0.266 | 7.034*** | -0.248(*) | -0.175 | 1.940 |
| | Bolting Rate | -0.174 | 0.196 | 1.727 | -0.047 | 0.176 | 34.43*** |
| B. Reproductive Traits | Total # of flowers | 1.26*** | 0.136 | 88.305*** | 0.558*** | 0.062 | 9.38** |
| | Seeds per fruit | 0.008 | -0.432* | 8.805*** | 0.063 | -0.123* | 6.59* |
| C. Size Traits | Height at first flower | -0.334 | 1.074*** | 97.61*** | -0.058 | 0.511*** | 3.83(*) |
| | Total Branches | 0.636** | 0.044 | 14.94*** | 0.116** | 0.121 | 8.27** |
| | Biomass | 1.10*** | 0.842*** | 133.78*** | 0.375*** | 0.634*** | 20.19*** |

Table 5A. ANOVA testing the effects of generation on expression of reproductive traits of *C. americanum* populations. The numerator degrees of freedom is one. F-ratio is listed for each effect.


| Trait | Site | Gen | Cross | Site x Gen | Cross x Gen | Cross x Site |
|--------------------|----------|---------|-------|------------|-------------|--------------|
| Total # of seeds | 19.65*** | 3.04(*) | 0.81 | 0.00 | 0.36 | 0.70 |
| Total # of fruit | 15.16*** | 2.32 | 0.47 | 0.04 | 0.31 | 0.99 |
| Total # of flowers | 9.98** | 1.16 | 1.85 | 0.32 | 0.42 | 0.78 |
| Biomass | 24.20*** | 7.56** | 2.64* | 1.97 | 0.84 | 3.28** |
| d.f numerator | 1 | 2 | 5 | 1 | 3 | 5 |

**d.f. denominator: 545-673

Table 5B. Linear Contrast of L.S means generated by ANOVAs for reproductive traits that quantifies the level of reproductive fitness exhibited by each generation (P, F2, RF2) of *C. americanum* populations. Reciprocal crosses were also included these analyzes and were noted by placing a capital R in front of the generations associated with the cross. P-values indicate whether F2 (F2 +RF2) means are significantly different from parents. The numerator d.f. is one. An arrow in the mean comparison column indicates if F2s had means lower or higher than the parents.

| Traits | d.f. denominator | F-ratio | p-value | Mean Comparison |
|-------------------------|------------------|---------|---------|-----------------|
| Number of Seeds/fruit | 547 | 8.83 | 0.003 | ↓ |
| Biomass | 555 | 7.94 | 0.005 | ↑ |
| Total number of fruits | 507 | 0.13 | 0.777 | = |
| Total number of flowers | 675 | 0.11 | 0.746 | = |

Figure 1. Location of nine *Campanulastrum americanum* populations (AL, NE, MI, TN, OK, MN, MS, WI, and OH) planted in common garden sites, Pierce Cedar Creek Institute in Hastings (MI) and CSU's Oxbow Meadows Environmental Learning Center in Columbus (GA). Common garden site locations are noted with stars. Red lines indicated which populations were crossed to generate F1s and F2s.






Figure 2. Comparing L.S. mean (\pm S.E.) A) minimum number of days to germination, B) percent germination across northern and southern *C. americanum* populations at Georgia and Michigan common garden sites. Means reflect germination data that was collected for a period of two years (2012-2014).






Figure 3. Assessing variations in phenology. Comparing log+1-transformed mean (\pm S.E.) A) bolting rate, B) number of days to first flower, and C) minimum number of days to fruit maturation across northern and southern *C. americanum* populations at Georgia and Michigan common garden sites (2014).






Figure 4. Assessing traits indicating size. Comparing log+1-transformed mean (\pm S.E.) A) height at flowering, B) total branch length, and C) biomass across northern and southern *C. americanum* populations at Georgia and Michigan common garden sites (2014).






Figure 5. Assessing variation in reproductive traits. Comparing log+1-transformed mean (\pm S.E.) A) total number of flowers, B) total number of fruits, C) seeds per fruit, and across northern and southern *C. americanum* populations at Georgia and Michigan common garden sites (2014).

Figure 6. Comparing log+1-transformed cumulative fitness means of source origins within each common garden site. Cumulative fitness was calculated by multiplying total number of fruits by number of seeds per fruit. Data collected from the rosette common garden experiment (2014) was used in this calculation.





Figure 7. Assessing hybrid breakdown by comparing L.S. means (\pm S.E.) A) total number of flowers, B) total number of fruits, C) number of seeds/fruit, and D) biomass. Linear contrasts were performed to detect significance between means of parents and F2s of *C. americanum* populations.



Appendix A: Geographical locations of nine *Campanulastrum americanum* populations (AL, NE, MI, TN, OK, MN, MS, WI, and OH) planted in common garden sites.

| Populations | Latitude | Longitude |
|-------------|----------|-----------|
| MS | 31.74 | -88.52 |
| AL | 31.92 | -86.69 |
| OK | 33.95 | -94.57 |
| TN | 35.77 | -88.06 |
| NE | 40.75 | -96.72 |
| OH | 41.12 | -81.52 |
| MI | 42.30 | -85.36 |
| WI | 43.35 | -89.95 |
| MN | 44.82 | -93.31 |

Appendix B1: Pearson product-moment correlation coefficients for reproductive and phenological traits of southern and northern *C. americanum* populations planted in the Georgia common garden site (2014).

| | | Number of Days | HAF (cm) | Total Branch Length (cm) | Seeds per fruit | Biomass (g) | Number of fruits | TOTAL FLOWERS | Ratio |
|--------------------------|---------------------|----------------|----------|--------------------------|-----------------|-------------|------------------|---------------|-------|
| Number of Days | Pearson Correlation | 1 | -.245** | -.153 | .081 | -.247** | -.267** | -.060 | .150 |
| | Sig. (2-tailed) | | .007 | .090 | .444 | .007 | .004 | .507 | .125 |
| | N | 123 | 122 | 123 | 92 | 119 | 112 | 123 | 106 |
| HAF (cm) | Pearson Correlation | -.245** | 1 | .263** | .271** | .467** | .526** | .534** | .013 |
| | Sig. (2-tailed) | .007 | | .003 | .009 | .000 | .000 | .000 | .898 |
| | N | 122 | 122 | 122 | 91 | 118 | 111 | 122 | 106 |
| Total Branch Length (cm) | Pearson Correlation | -.153 | .263** | 1 | .072 | .086 | .455** | .406** | -.144 |
| | Sig. (2-tailed) | .090 | .003 | | .470 | .334 | .000 | .000 | .142 |
| | N | 123 | 122 | 155 | 104 | 127 | 113 | 155 | 106 |
| Seeds per fruit | Pearson Correlation | .081 | .271** | .072 | 1 | .268** | .308** | .300** | -.135 |
| | Sig. (2-tailed) | .444 | .009 | .470 | | .009 | .004 | .002 | .233 |
| | N | 92 | 91 | 104 | 104 | 95 | 88 | 104 | 80 |
| Biomass (g) | Pearson Correlation | -.247** | .467** | .086 | .268** | 1 | .685** | .488** | .043 |
| | Sig. (2-tailed) | .007 | .000 | .334 | .009 | | .000 | .000 | .664 |
| | N | 119 | 118 | 127 | 95 | 127 | 112 | 127 | 103 |
| Number of fruits | Pearson Correlation | -.267** | .526** | .455** | .308** | .685** | 1 | .812** | -.138 |
| | Sig. (2-tailed) | .004 | .000 | .000 | .004 | .000 | | .000 | .172 |
| | N | 112 | 111 | 113 | 88 | 112 | 113 | 113 | 99 |
| TOTAL FLOWERS | Pearson Correlation | -.060 | .534** | .406** | .300** | .488** | .812** | 1 | -.108 |
| | Sig. (2-tailed) | .507 | .000 | .000 | .002 | .000 | .000 | | .270 |
| | N | 123 | 122 | 155 | 104 | 127 | 113 | 155 | 106 |
| Ratio | Pearson Correlation | .150 | .013 | -.144 | -.135 | .043 | -.138 | -.108 | 1 |
| | Sig. (2-tailed) | .125 | .898 | .142 | .233 | .664 | .172 | .270 | |
| | N | 106 | 106 | 106 | 80 | 103 | 99 | 106 | 106 |

** . Correlation is significant at the 0.01 level (2-tailed).

Appendix B2: Pearson product-moment correlation coefficients for reproductive and phenological traits of southern and northern *C. americanum* populations planted in the Michigan common garden site (2014).

| | | Number of Days | HAF (cm) | Total Branch Length (cm) | Seeds per fruit | Biomass (g) | Number of fruits | TOTAL FLOWERS | Ratio |
|--------------------------|---------------------|----------------|----------|--------------------------|-----------------|-------------|------------------|---------------|--------|
| Number of Days | Pearson Correlation | 1 | -.323** | -.024 | -.079 | -.172 | -.265** | -.155 | .390** |
| | Sig. (2-tailed) | | .000 | .789 | .375 | .052 | .003 | .080 | .000 |
| | N | 128 | 128 | 128 | 128 | 128 | 127 | 128 | 128 |
| HAF (cm) | Pearson Correlation | -.323** | 1 | .112 | .298** | .814** | .867** | .381** | -.078 |
| | Sig. (2-tailed) | .000 | | .205 | .001 | .000 | .000 | .000 | .381 |
| | N | 128 | 129 | 129 | 129 | 129 | 128 | 129 | 129 |
| Total Branch Length (cm) | Pearson Correlation | -.024 | .112 | 1 | .071 | .155* | .130 | -.040 | .049 |
| | Sig. (2-tailed) | .789 | .205 | | .324 | .041 | .099 | .573 | .579 |
| | N | 128 | 129 | 203 | 195 | 174 | 162 | 203 | 129 |
| Seeds per fruit | Pearson Correlation | -.079 | .298** | .071 | 1 | .194* | .228** | .218** | .124 |
| | Sig. (2-tailed) | .375 | .001 | .324 | | .011 | .004 | .002 | .161 |
| | N | 128 | 129 | 195 | 195 | 171 | 161 | 195 | 129 |
| Biomass (g) | Pearson Correlation | -.172 | .814** | .155* | .194* | 1 | .751** | .418** | -.014 |
| | Sig. (2-tailed) | .052 | .000 | .041 | .011 | | .000 | .000 | .875 |
| | N | 128 | 129 | 174 | 171 | 174 | 162 | 174 | 129 |
| Number of fruits | Pearson Correlation | -.265** | .867** | .130 | .228** | .751** | 1 | .416** | -.048 |
| | Sig. (2-tailed) | .003 | .000 | .099 | .004 | .000 | | .000 | .592 |
| | N | 127 | 128 | 162 | 161 | 162 | 162 | 162 | 128 |
| TOTAL FLOWERS | Pearson Correlation | -.155 | .381** | -.040 | .218** | .418** | .416** | 1 | .018 |
| | Sig. (2-tailed) | .080 | .000 | .573 | .002 | .000 | .000 | | .840 |
| | N | 128 | 129 | 203 | 195 | 174 | 162 | 203 | 129 |
| Ratio | Pearson Correlation | .390** | -.078 | .049 | .124 | -.014 | -.048 | .018 | 1 |
| | Sig. (2-tailed) | .000 | .381 | .579 | .161 | .875 | .592 | .840 | |
| | N | 128 | 129 | 129 | 129 | 129 | 128 | 129 | 129 |

** . Correlation is significant at the 0.01 level (2-tailed).

* . Correlation is significant at the 0.05 level (2-tailed).





Appendix C: Detecting hybrid breakdown by analyzing whether a linear relationship $((P1+P2)/2=(F2+rF2)/2)$ exists across two generations of *C. americanum* (P, F2/RF2) for each population cross-type. Generations associated with reciprocal crosses were noted by placing R in front of the generation category. Contrast linear analyses were performed on a selected group of reproductive traits that assess the level of reproductive fitness exhibited by each generation at the Georgia common garden site. The numerator d.f. is one. Arrows in the mean comparison column indicates if F2s had means lower or higher than the parents.

| Population cross-type | Trait | d.f Error | F-value | p-value | Mean comparison |
|-----------------------|-------------------------|-----------|---------|---------|-----------------|
| AL69 x NE59 | Total number of seeds | 31 | 11.24 | 0.0021 | |
| | Biomass | 43 | 3.67 | 0.0624 | = |
| | Total number of fruits | 50 | 2.20 | 0.1440 | = |
| | Total number of flowers | 45 | 1.80 | 0.1870 | = |
| MI44 x MS55 | Total number of seeds | 39 | 2.21 | 0.1232 | = |
| | Biomass | 51 | 8.75 | 0.0047 | |
| | Total number of fruits | 45 | 1.12 | 0.2948 | = |
| | Total number of flowers | 43 | 0.33 | 0.5655 | = |
| OK61 x MN38 | Total number of seeds | 26 | 1.37 | 0.1395 | = |
| | Biomass | 34 | 2.11 | 0.0652 | = |
| | Total number of fruits | 29 | 0.68 | 0.4097 | = |
| | Total number of flowers | 34 | 2.13 | 0.1536 | = |
| MS55 X WI14 | Total number of seeds | 43 | 7.47 | 0.0091 | |
| | Biomass | 58 | 2.01 | 0.1617 | = |
| | Total number of fruits | 45 | 1.58 | 0.2154 | = |
| | Total number of flowers | 46 | 1.40 | 0.2424 | = |
| OH64 X OK61 | Total number of seeds | 36 | 0.00 | 0.9941 | = |
| | Biomass | 43 | 0.02 | 0.8573 | = |
| | Total number of fruits | 37 | 0.00 | 0.9590 | = |
| | Total number of flowers | 42 | 1.61 | 0.2113 | = |
| MN38 X TN19 | Total number of seeds | 22 | 3.89 | 0.0613 | = |
| | Biomass | 30 | 0.33 | 0.5707 | = |
| | Total number of fruits | 23 | 0.45 | 0.5097 | = |
| | Total number of flowers | 29 | 0.02 | 0.8990 | = |





Appendix D: Detecting hybrid breakdown by analyzing whether a linear relationship $((P1+P2)/2=(F2+rF2)/2)$ exists across two generations of *C. americanum* (P, F2/RF2) for each population cross-type. Generations associated with reciprocal crosses were noted by placing R in front of the generation category. Contrast linear analyzes were performed on a selected group of reproductive traits that assess the level of reproductive fitness exhibited by each generation at the Michigan common garden site. The numerator d.f. is one. Arrows in the mean comparison column indicates if F2s had means lower or higher than the parents.

| Population cross-type | Trait | d.f Error | F-value | p-value | Mean Comparison |
|-----------------------|-------------------------|-----------|---------|---------|-----------------|
| AL69 x NE59 | Total number of seeds | 36 | 0.04 | 0.8401 | = |
| | Biomass | 54 | 0.06 | 0.8034 | = |
| | Total number of fruits | 50 | 2.48 | 0.1215 | = |
| | Total number of flowers | 59 | 4.75 | 0.0332 | = |
| MI44 x MS55 | Total number of seeds | 66 | 0.00 | 0.9888 | = |
| | Biomass | 82 | 2.49 | 0.1187 | = |
| | Total number of fruits | 72 | 5.39 | 0.0231 | = |
| | Total number of flowers | 73 | 0.24 | 0.6260 | = |
| OK61 x MN38 | Total number of seeds | 39 | 0.07 | 0.8001 | = |
| | Biomass | 44 | 0.17 | 0.6808 | = |
| | Total number of fruits | 39 | 0.29 | 0.5939 | = |
| | Total number of flowers | 42 | 4.70 | 0.0359 | = |
| MS55 X WI14 | Total number of seeds | 53 | 1.47 | 0.3155 | = |
| | Biomass | 70 | 3.15 | 0.0805 | = |
| | Total number of fruits | 61 | 3.20 | 0.0787 | = |
| | Total number of flowers | 64 | 0.13 | 0.7178 | = |
| OH64 X OK61 | Total number of seeds | 59 | 0.15 | 0.6977 | = |
| | Biomass | 58 | 0.04 | 0.8334 | = |
| | Total number of fruits | 77 | 0.33 | 0.5660 | = |
| | Total number of flowers | 63 | 0.39 | 0.5371 | = |
| MN38 X TN19 | Total number of seeds | 54 | 0.17 | 0.6788 | = |
| | Biomass | 58 | 0.64 | 0.4283 | = |
| | Total number of fruits | 52 | 0.20 | 0.6562 | = |
| | Total number of flowers | 54 | 0.01 | 0.9259 | = |

Appendix E: Assessing variations in phenology across *C. americanum* populations. Comparing L.S. means A) number of days to germination, B) bolting rate, C) number of days to first flower, and D) number of days to fruit maturation across northern and southern populations at Georgia and Michigan common garden sites (2014).



Appendix F: Assessing variations in reproductive fitness across *C. americanum* populations. Comparing L.S. means A) percent germination, B) total number of flowers, C) number of fruits, and D) number of seeds per fruit across northern and southern populations at Georgia and Michigan common garden sites (germ data from 2012-2013; rosettes during 2014).



Appendix G: Assessing variations in size across *C. americanum* populations. Comparing L.S. means A) total branch length, B) biomass, and C) height at flowering across northern and southern populations at Georgia and Michigan common garden sites (2014).

